



INTERNATIONAL  
BIRD RESCUE



## DID YOU KNOW?

Wildlife rehabilitators aren't the only ones who can protect birds from bird flu, **YOU** can make a difference too!



### DISINFECT YOUR SHOES

Be sure to disinfect the bottom of your shoes as bird flu can be spread through bird droppings. Use diluted bleach and leave shoes in the sun to dry.



### MANAGE BIRDFEEDERS RESPONSIBLY

Although songbirds and hummingbirds have a lower risk, birdfeeders and bird baths should be cleaned and disinfected at least weekly.



### REPORT SICK OR DECEASED BIRDS

For birds experiencing symptoms like tremors, bluish eyes, head-twisting, seizures, or respiratory problems, alert your state's wildlife agency. Otherwise, contact your local wildlife rehabilitator before intervening.



### BE AWARE OF ASYMPTOMATIC SPECIES

Mallards, ducklings, and others may be infected but not show symptoms of the disease.



### TAKE SPECIAL CARE AROUND DOMESTIC BIRDS

If you are around chickens, ducks, and turkeys—use gloves, change clothing and shoes to avoid cross-contamination.



### SUPPORT YOUR LOCAL WILDLIFE REHABILITATOR

Wildlife organizations cannot do it alone. Support your local group with time and resources to help stop the spread of this virus.